

Zinc Extra

Contains: 15% w/v elemental Zinc and 4% w/v elemental Nitrogen (as Nitrate N)

An Inorganic Micronutrient for foliar application. Prevents & corrects Zinc deficiency in all arable crops.

Rates of use:

Deficiency Status:	Application Rate:
Maintenance	2.0 litres/ha
Moderate	4.0 litres/ha
Severe	4.0 litres/ha & repeat

Water Volumes:

Conventional sprayer : 200 L/ha minimum
Low volume sprayer : 55 L/ha minimum

Crop recommendations & growth stages:

CROP	TIMING
Cereals	6 tillers to 2nd node
Maize	6 to 8 leaves
Top Fruit	After petal fall
Potatoes	From 7 days after 100% emergence to when rows meet.
Vegetables	When sufficient leaf area
Other Crops	When sufficient leaf area



DE SANGOSSE

De Sangosse Ltd.
Swaffham Bulbeck,
Cambridge,
CB25 0LU
Tel: (01223) 811215
Fax: (01223) 810020
email: info@desangosse.co.uk
web: www.desangosse.co.uk

Contents:
10 Litres e

L178A

UN No. 3082:
Environmentally Hazardous
Substance, Liquid, NOS

Precautions:

Harmful if swallowed.

Irritating to eyes, respiratory system and skin.


Very toxic to aquatic organisms, may cause long-term adverse effects in the aquatic environment.

Do not breathe fumes. Avoid contact with skin and eyes. In case of contact with eyes, rinse immediately with plenty of water and seek medical advice. After contact with skin, wash immediately with plenty of water. Wear suitable protective clothing, gloves and eye / face protection. This material and its container must be disposed of as hazardous waste. Avoid release to the environment. Refer to special instructions / safety data sheets.

PROTECT FROM FROST.



HARMFUL



DANGEROUS For
The ENVIRONMENT