



**DE SANGOSSE**

## Spray Enhancers

### Phase-II



#### What is it?

Phase-II is a high quality new generation methylated seed oil (MSO) adjuvant oil derived from oilseed rape.

#### ADJ No.

0622

PCS No. 92147

#### Formulation

An emulsifiable concentrate containing 95% w/w (842 g/L) esterified rapeseed oil.

#### Main Recommendations

Phase-II may be used at a maximum concentration of 1.78% spray solution (maximum 2 litres per hectare) on the following crops, up to the growth stages specified, with all approved pesticides applied up to their full approved rate.

Crop	Growth stages before which Phase-II may be used with approved pesticides
Cereals	Up to and including GS 52 (1/4 inflorescence)
Oilseed rape	Up to and including 10% potential pods
Broad, dwarf French, field & runner bean	Up to and including first pod set
Combining & vining pea	Up to and including flat pod (non-edible pods only)
Edible podded pea	Up to and including pod set
Linseed	Up to and including 10% capsules formed
Forage maize	Up to and including tip of tassel visible
Sugar beet	Up to and including 6 fully expanded leaves
Potato	Up to and including tuber initiation
Brussels sprout	Up to and including lateral buds begin to develop
Cauliflower	Up to and including cauliflower heads begin to form; width of growing tip up to 1 cm <sup>2</sup>

Phase-II can be used with all approved pesticides on all edible crops provided that the statutory conditions of use for the pesticide are followed and that the pesticide is not used at more than 50% of the maximum approved rate for that application. It can be used with all approved pesticides on all non-edible crops provided that the statutory conditions of use for the pesticide are followed.

Phase-II can be used as an alternative to a mineral oil.

Phase-II can be used to reduce drift.

#### Rate of Use

This adjuvant may be used at a maximum concentration of 1.78% spray solution. Phase-II is normally used at 0.5% dilution of the final water volume, minimum 500 ml per hectare.

#### Water Volume

The water volume is governed by that stated on the partner pesticide label.





**DE SANGOSSE**

## Spray Enhancers

### Phase-II



#### Mixing Sequence

Phase-II is compatible in all situations where an adjuvant oil is recommended or required. Three quarter fill the spray tank with water and commence agitation. Add the required quantity of Phase-II. Add pesticide(s) as directed. Continue agitation while topping up the tank and during spraying.

#### Pack Size

5 litres

